

MYCOTAXON

<http://dx.doi.org/10.5248/127.59>

Volume 127, pp. 59–62

January–March 2014

***Diplococcium variegatum*, a new conidial fungus from the semi-arid Caatinga biome of Brazil**

SILVANA SANTOS DA SILVA¹, ALISSON CARDOSO RODRIGUES DA CRUZ¹,
LUÍS FERNANDO PASCHOLATI GUSMÃO^{1*}, & RAFAEL F. CASTAÑEDA-RUIZ²

¹Universidade Estadual de Feira de Santana,

Departamento de Ciências Biológicas, Laboratório de Micologia,

Avenida Transnordestina, s/n, Novo Horizonte, 44036-900, Feira de Santana, BA, Brazil.

²Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt'

(INIFAT), Académico Titular de la Academia de Ciencias de Cuba,

Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

*CORRESPONDENCE TO: lgusmao@uefs.br

ABSTRACT — *Diplococcium variegatum* sp. nov., found on a decaying leaf of an unidentified plant, is described and illustrated. It is characterized by variegated pigmented structures and limoniform to ovoid, 0-septate, pale brown, brown to dark brown conidia in acropetal chains.

KEY WORDS — hyphomycetes, plant debris, taxonomy

Introduction

During surveys for conidial fungi on plant debris in a semi-arid region of northeast Brazil, an interesting specimen of *Diplococcium* Grove was found. The fungus clearly differs morphologically from all described species and is therefore described as new.

Materials & methods

Expeditions to Serra da Jibóia, Bahia, Brazil were made every 3 months, from October 2008 to April 2009. Samples of plant debris (twigs, bark, leaves, and petioles) were collected in paper bags. In the laboratory the samples were placed in Petri dish moist chambers and stored in a 170 L polystyrene box with 200 mL sterile water plus 2 mL glycerol at 25°C for 30 days (Castañeda-Ruiz 2005). During this period, the samples were examined for the presence of conidial fungi. Slides were prepared in polyvinyl alcohol plus lactic acid and phenol. Measurements were made at a magnification of ×1000. Micrographs were obtained with an Olympus BX 51.

Taxonomy

Diplococcium variegatum S.S. Silva, Gusmão & R.F. Castañeda, **sp. nov.** FIG. 1

MYCOBANK MB 805030

Differs from *Diplococcium parcum* by larger limoniform to ovoid conidia and the variably colored conidiogenous cells.

TYPE: Brazil. Bahia State: Santa Teresinha, Serra da Jibóia, on decaying leaves of an unidentified plant, 19.II.2009; coll. S.S. Silva. (Holotype: HUEFS 194254).

ETYMOLOGY: Latin, *variegatum*, referring to the variable coloration of the conidiogenous cells and conidia.

COLONIES effuse, hairy, brown. Mycelium superficial and immersed in substrate. CONIDIOPHORES macronematous, mononematous, erect, straight to slightly flexuous, simple or branched, 4–7-septate, smooth, brown below, variegated pigmented towards the apex, $135\text{--}189 \times 12.0\text{--}16.5\ \mu\text{m}$, cell basal lobed $21\text{--}45\ \mu\text{m}$ wide. CONIDIOGENOUS CELLS polytretic, integrated, terminal and intercalary, pale brown to dark brown to reddish brown, $19.5\text{--}25 \times 5.0\text{--}6.5\ \mu\text{m}$. CONIDIA limoniform to ovoid, unicellular, smooth, thick-walled, pale brown to dark brown or reddish brown, $7.5\text{--}22.5 \times 5\text{--}12\ \mu\text{m}$, forming acropetal chains. Teleomorph unknown.

NOTE: *Diplococcium* was described by Grove (1885) with the type species *D. spicatum* Grove from dead wood in England. Braun et al. (1996), Castañeda-Ruiz & Kendrick (1991), Cruz et al. (2007), Goh & Hyde (1998), Hernández-Restrepo et al. (2012), Ma et al. (2012), Pirozynski (1972), and Wang & Sutton (1998) documented 28 accepted species of *Diplococcium*, which are characterized by terminal or intercalary polytretic conidiogenous cells that produce 0–7-septate conidia in acropetal chains. *Diplococcium capitatum* Piroz., *D. hughesii* C.J.K. Wang & B. Sutton, and *D. parcum* Hol.-Jech. are also species that produce exclusively aseptate conidia. *Diplococcium capitatum* differs from *D. variegatum* by conidiogenous pores confined to a swollen terminal cell that produces short cylindric or broadly ellipsoid conidia (Pirozynski 1972); *D. hughesii* is separated by its subglobose oval to oblong smaller ($7\text{--}12 \times 5\text{--}7\ \mu\text{m}$) conidia and highly branched longer conidiophores (Holubová-Jechová 1982, Wang & Sutton 1998), while *D. parcum* produces smaller ($8.5\text{--}13 \times 5.5\text{--}8\ \mu\text{m}$) ovate to obpyriform or ellipsoidal conidia (Holubová-Jechová 1982).

Acknowledgments

The authors express their sincere gratitude to Dr. De-Wei Li and Prof. Dr. Bryce Kendrick for their critical review of the manuscript. The authors thank the Program of Research on Biodiversity in the Brazilian semi-arid (PPBIO semi-arid/ Ministry of Technology and Science). RFCR is grateful to the Cuban Ministry of Agriculture and “Programa de Salud Animal y Vegetal,” project P131LH003033 Cuban Ministry of Science, Technology and Environment for facilities.



FIG 1. *Diplococcium variegatum* (ex HUEFS 194254). A. Conidiophore and conidia. B. Conidiogenous cell and pores (indicated by arrows). C. Conidiogenous cells and conidia. D–F. Conidia. Scale bars: A = 20 μ m; B–F = 5 μ m.

Literature cited

- Braun U, Hosagoudar VB, Abraham TK. 1996. *Diplococcium atrovelutinum* sp. nov. from India. New Botanist 23: 1–4.
- Castañeda-Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. Anais do V Congresso Latino Americano de Micología, Brasília: 182–183.

- Castañeda-Ruiz RF, Kendrick B. 1991. Ninety conidial fungi from Cuba and three from Canada. University of Waterloo, Biology Series 35: 1–132.
- Cruz ACR, Gusmão LFP, Leão-Ferreira SM, Castañeda-Ruiz RF. 2007. Conidial fungi from the semi-arid Caatinga biome of Brazil. *Diplococcium verruculosum* sp. nov. and *Lobatopedis longirostratum* sp. nov. Mycotaxon 103: 33–38.
- Goh TK, Hyde KD. 1998. A synopsis of and a key to *Diplococcium* species, based on the literature, with a description of a new species. Fungal Diversity 1: 65–83.
- Grove WB. 1885. New or noteworthy fungi. Part II. J. Bot. 23: 161–169.
- Hernández-Restrepo M, Silvera-Simón C, Mena-Portales J, Mercado-Sierra Á, Guarro J, Gené J. 2012. Three new species and a new record of *Diplococcium* from plant debris in Spain. Mycol. Prog. 11: 191–199. <http://dx.doi.org/10.1007/s11557-011-0741-6>
- Holubová-Jechová V. 1982. Lignicolous hyphomycetes from Czechoslovakia 6. *Spadicoides* and *Diplococcium*. Folia Geobot. Phytotax. 17: 295–327.
- Ma LG, Ma J, Zhang YD, Zhang XG. 2012. *Spadicoides camelliae* and *Diplococcium livistonae*, two new hyphomycetes on dead branches from Fujian Province, China. Mycoscience 53: 25–30. <http://dx.doi.org/10.1007/s10267-011-0138-z>
- Pirozynski KA. 1972. Microfungi of Tanzania. I. Miscellaneous fungi on oil palm. II. New hyphomycetes. Mycol. Pap. 129: 1–64.
- Wang CJK, Sutton B. 1998. *Diplococcium hughesii* sp. nov. with a *Selenosporella* synanamoph. Can. J. Bot. 76: 1608–1613. <http://dx.doi.org/10.1139/b98-105>